

## Appendix A: Birds

### Blue-winged Warbler

*Vermivora cyanoptera*

Federal Listing	N/A
State Listing	N/A
Global Rank	G5
State Rank	S4
Regional Status	Very High



Photo by USFWS

#### Justification (Reason for Concern in NH)

Populations of many shrubland birds are in strong decline, both in the Northeast and sometimes across larger portions of their continental ranges. For this reason, most species were included in the Northeast list of SGCN, with those that occur regularly in NH retained for the NH WAP revision. Across its range as a whole, the Blue-winged Warbler has been experiencing non-significant declines of less than 1%/year since 1966 (Sauer et al. 2014). Declines appear strongest in the east, and in BCR 30 (where most of NH's Blue-winged Warblers occur) are significantly negative (roughly -2%/year). Repeated Breeding Bird Atlases in the northeast have documented range expansion to the north (McGowan and Corwin 2008, Renfrew 2013), or filling in of previously unoccupied areas (MassAudubon 2014). In New Hampshire, there are not enough BBS data to determine a trend, but the species is now found farther north and west than was documented during the NH Atlas in the early 1980s (Foss 1994).

#### Distribution

The Blue-winged Warbler breeds in the eastern U.S. from Minnesota and Maine south to Arkansas and Georgia. It winters along the Caribbean slope of Central America from southern Mexico to Panama (Gill et al. 2001). In New Hampshire it occurs primarily south and east of Concord of the White Mountains, with scattered records in the southern Connecticut River Valley. Within this distribution it is very locally distributed, with most records coming from southern Strafford and eastern Rockingham counties.

#### Habitat

Like all shrubland birds, this species occurs in habitats dominated by shrubs or young trees, sometimes interspersed with mature trees (e.g., pine barrens) or open bare or grassy areas. Typical examples in New Hampshire include regenerating timber harvests, power line rights-of-way, shrubby old fields and edges, and pine barrens. From a bird perspective, such habitats can be subdivided into those dominated by shrubs vs. dominated by saplings. The former – sometimes referred to as “scrub-shrub” – is more typical of abandoned old fields, utility rights-of-way, and open areas within pine barrens. Such habitats often persist for relatively long periods without the need for additional management. Saplings, on the other hand, are typical of areas subject to timber harvest, and rarely retain early successional characteristics beyond 15-20 years. These are also regularly referred to as “young forest.” Prairie Warblers are rare in the latter habitat, and tend to reach their highest densities in pine barrens or other areas with scattered short conifers (Hunt 2013).


## Appendix A: Birds

### NH Wildlife Action Plan Habitats

- Shrublands
- Pine Barrens

Distribution of  
BLUE-WINGED WARBLER  
in New Hampshire

■ Current Range  
▨ Localized



Distribution Map

### Current Species and Habitat Condition in New Hampshire

Significant population declines rangewide, but increases and northward range expansion at the northern edge of its range (see Justification).

### Population Management Status

Management is not currently in place for this species.

### Regulatory Protection (for explanations, see Appendix I)

- Migratory Bird Treaty Act (1918)

### Quality of Habitat

Highly variable – see shrubland and pine barrens habitat profiles

### Habitat Protection Status

Highly variable – see shrubland and pine barrens habitat profiles

### Habitat Management Status

Habitat management has not been implemented specifically for this species, although management does occur for other species (American Woodcock, New England Cottontail) that often use the same habitats. See also shrubland and pine barrens habitat profiles.

## Appendix A: Birds

### Threats to this Species or Habitat in NH

*Threat rankings were calculated by groups of taxonomic or habitat experts using a multistep process (details in Chapter 4). Each threat was ranked for these factors: Spatial Extent, Severity, Immediacy, Certainty, and Reversibility (ability to address the threat). These combined scores produced one overall threat score. Only threats that received a “medium” or “high” score have accompanying text in this profile. Threats that have a low spatial extent, are unlikely to occur in the next ten years, or there is uncertainty in the data will be ranked lower due to these factors.*

#### **Habitat conversion due to development and impacts from fragmentation (Threat Rank: High)**

Ongoing residential and commercial development results in permanent loss of habitats for wildlife. Because many of the habitats used by shrubland birds are already embedded in developed landscapes (e.g., right-of-way, old fields) or viewed as “undesirable” or “waste” habitats, they may be more vulnerable to this threat.

#### **Habitat degradation and conversion from natural succession or a lack of active management (Threat Rank: High)**

In the absence of disturbance or management, the early successional and edge habitats preferred by this species generally revert to closed forest systems that are not heavily used, and as a result forest maturation is generally considered the most significant threat facing birds that use shrublands and young forests. See shrubland habitat profile for more information.

#### **Habitat and species impacts from invasive or introduced plants (Threat Rank: Medium)**

Non-native plants, particularly shrubs, have been demonstrated to have several negative effects on birds using shrubland habitats. Insect prey (particularly caterpillars) are usually less common on non-native shrubs (Burghardt et al. 2008, Fickenscher et al. 2014), while data on the nutritional value of fruit are more equivocal (e.g., Davis 2011). In some cases, birds experience lower reproductive success in non-native shrubs, although there is considerable variation (Rodewald et al. 2010, Schlossberg and King 2010), and local predator communities play an important role as well. In all cases, the effects of invasives on shrubland birds depend to a large extent on their relative abundance. If plant diversity is high, the negative effects are diluted and less likely to impact bird populations. However, if the habitat tends toward a monoculture, reduced insect supplies and/or higher predation may reduce reproductive success to the extent that the habitat becomes a sink.

#### **Mortality from subsidized or introduced predators (Threat Rank: Medium)**

Many predators (e.g., skunks, raccoons, feral cats) occur in relatively high densities in developed landscapes, often because of direct association with humans or food that is provided either intentionally or unintentionally. Most early successional birds nest on or near the ground, and as a result are more susceptible to nest predation. The problem is compounded because much early successional habitat is near human population centers.

#### **Habitat degradation from aspects of right-of-way management (Threat Rank: Medium)**

Rights-of-way need to be maintained as short vegetation so as to reduce risks associated with trees and powerlines. As a result these corridors are regularly treated by mechanical (rarely chemical) means to remove or cut back vegetation. In general, such practices create habitat suitable for shrubland birds, although in extreme cases a site may be rendered unsuitable for 1-2 years large

## ***Appendix A: Birds***

areas of vegetation are completely removed. If management occurs during the breeding season, reproductive success will be reduced. See also shrubland habitat profile.

### **List of Lower Ranking Threats:**

Habitat degradation from fire suppression and associated succession

Habitat degradation from sand and gravel pit reclamation practices that make habitat unsuitable

### **Actions to benefit this Species or Habitat in NH**

#### **Research on response to management**

**Primary Threat Addressed:** Habitat degradation and conversion from natural succession or a lack of active management

**Specific Threat (IUCN Threat Levels):** Natural system modifications

#### **Objective:**

To best conserve this and other shrubland birds, there is a recognized need for information on how it responds to management implemented for other early successional species.

#### **General Strategy:**

Collect standardize data on occupancy and abundance of Blue-winged Warblers at sites managed for New England Cottontail, American Woodcock, Karner Blue Butterfly, and other species. Combine these data with data on habitat availability at the state and regional scale, to estimate actual or potential population size, which in turn could be compared to population objectives that also need to be developed.

#### **Political Location:**

Statewide

#### **Watershed Location:**

Statewide

### **References, Data Sources and Authors**

#### **Data Sources**

Trend data from Breeding Bird Survey (Sauer et al. 2014, above). NH distribution data from NHBR/NH eBird

#### **Data Quality**

Because this species is easily detected and identifiable, data on distribution and habitat use are generally well known.

#### **2015 Authors:**

Pamela Hunt, NHA

#### **2005 Authors:**

## ***Appendix A: Birds***

### **Literature**

- Burghardt, K. T., D. W. Tallamy, and W. G. Shriver. 2008. Impact of native plants on bird and butterfly biodiversity in suburban landscapes. *Conservation Biology* 23:219-224.
- Davis, M. 2011. Do native birds care whether their berries are native or exotic? *No. Bioscience* 61:501-502.
- Fickenscher, J. L., J. A. Litvaitis, T. D. Lee, and P. C. Johnson. 2014. Insect responses to invasive shrubs: implications to managing thicket habitats in the northeastern United States. *Forest Ecology and Management*. 322:127-135.
- Foss, C.R. 1994. *Atlas of Breeding Birds in New Hampshire*. New Hampshire Audubon, Concord, NH.
- Gill, F.B., R.A. Canterbury and J.L. Confer. 2001. Blue-winged Warbler (*Vermivora cyanoptera*), *The Birds of North America Online* (A. Poole, Ed.). Ithaca: Cornell
- Massachusetts Audubon Society. 2014. Massachusetts Breeding Bird Atlas. Online results at: <http://www.massaudubon.org/our-conservation-work/wildlife-research-conservation/statewide-bird-monitoring/breeding-bird-atlases/bba2/>
- McGowan, K.J., and K. Corwin (eds.). 2008. *The Second Atlas of Breeding Birds in New York State*. Cornell University Press, Ithaca, NY.
- Renfrew, R.B. 2013 (ed.). *The Second Atlas of Breeding Birds in Vermont*. University Press of New England, Hanover, NH.
- Rodewald, A. D., D. P. Shustack, and L. E. Hitchcock. 2010. Exotic shrubs as ephemeral ecological traps for nesting birds. *Biological Invasions* 12:33-39.
- Sauer, J.R., J.E. Hines, J.E. Fallon, K.L. Pardieck, D.J. Ziolkowski, Jr., and W.A. Link. 2014. *The North American Breeding Bird Survey, Results and Analysis 1966 - 2013*. Version
- Schlossberg, S., and D. I. King. 2010. Effects of invasive woody plants on avian nest site selection and nesting success in shrublands. *Animal Conservation* 13:286-293.